



## AP<sup>®</sup> Physics C 1998 Scoring Guidelines

**The materials included in these files are intended for use by AP teachers for course and exam preparation in the classroom; permission for any other use must be sought from the Advanced Placement Program<sup>®</sup>. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service<sup>®</sup> (ETS<sup>®</sup>), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 4,200 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2002 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, SAT, and the acorn logo are registered trademarks of the College Entrance Examination Board. APIEL is a trademark owned by the College Entrance Examination Board. PSAT/NMSQT is a registered trademark jointly owned by the College Entrance Examination Board and the National Merit Scholarship Corporation. Educational Testing Service and ETS are registered trademarks of Educational Testing Service.

1998 Physics C Solutions

Distribution  
of points

Mech. 1 (15 points)

(a)

i. 1 point

For correct answer

1 point

$$\bar{v} = \frac{\Delta s}{\Delta t} = \frac{0.30 \text{ m} - 0.10 \text{ m}}{0.30 \text{ s} - 0.10 \text{ s}} = 1.00 \text{ m/s}$$

ii. 1 point

For correct answer

1 point

$$\bar{v} = \frac{\Delta s}{\Delta t} = \frac{0.99 \text{ m} - 0.87 \text{ m}}{1.10 \text{ s} - 0.90 \text{ s}} = 0.60 \text{ m/s}$$


iii. 1 point

For correct answer

1 point

$$\bar{v} = \frac{\Delta s}{\Delta t} = \frac{1.18 \text{ m} - 1.14 \text{ m}}{1.90 \text{ s} - 1.70 \text{ s}} = 0.20 \text{ m/s}$$

(b) 3 points



For line I horizontal at  $v = 1.00$  m/s or at answer obtained for (a)i.

1 point

For line II with monotonic, negative slope between points  $P_1$  and  $P_2$ ,

$P_1$  at (0.70 - 0.80, 1.00) or (0.70 - 0.80, answer to (a)i.), and

$P_2$  at (1.20 - 1.30, 0.20) or (1.20 - 1.30, answer to (a)iii.)

1 point

(This line may be straight, which is consistent with the data, or slightly curved as shown, which is consistent with the behavior of springs)

For line III horizontal at  $v = 0.20$  m/s or at answer obtained for (a)iii.

1 point

Mech. 1 (continued)

(c)

i. 3 points

For any statement of conservation of momentum or energy

1 point

For proper conservation of momentum or energy equation

1 point

Method 1: Conservation of momentum

$$m_A v_{Ai} = m_A v_{Af} + m_B v_B$$

$$(0.90 \text{ kg})(1.00 \text{ m/s}) = (0.90 \text{ kg})(0.20 \text{ m/s}) + (0.60 \text{ kg})v_B$$

Method 2: Recognize from part (d) that energy is also conserved.


$$\frac{1}{2} m_A v_{Ai}^2 = \frac{1}{2} m_A v_{Af}^2 + \frac{1}{2} m_B v_B^2$$

$$\frac{1}{2} (0.90 \text{ kg})(1.00 \text{ m/s})^2 = \frac{1}{2} (0.90 \text{ kg})(0.20 \text{ m/s})^2 + \frac{1}{2} (0.60 \text{ kg})v_B^2$$

For correct answer

1 point

$$v_B = 1.2 \text{ m/s}$$

For line I horizontal at  $v = 0$ 

1 point

For line II with monotonic, positive slope between points  $P_1$  and  $P_2$ , $P_1$  at (0.70 - 0.80, 0), and $P_2$  at (1.20 - 1.30, 1.20) or (1.20 - 1.30, answer to (c)i.)

1 point

(This line may be straight, which is consistent with the data, or slightly curved as shown, which is consistent with the behavior of springs)

For line III horizontal at  $v = 1.20 \text{ m/s}$  or at answer obtained for (c)i.

1 point

## Mech. 1 (continued)

(d)

i. 2 points

For correct answer

1 point

Yes, the collision is elastic.

For any reasonable justification

1 point

Examples:

The final kinetic energy equals the initial kinetic energy.

The spring force is conservative meaning the total energy stored equals the total energy released.

The compressed spring stores and releases energy in equal amounts.

(The justification point was not awarded if student answered "no" to the question.)

ii. 1 point

For any reasonable explanation

1 point

Examples:

The compressed spring stores maximum amount of kinetic energy.

At time  $t = 1$  s, there is maximum kinetic energy stored as potential energy.At time  $t = 1$  s, the spring has maximum potential or elastic energy.

## 1998 Physics C Solutions

Distribution  
of points

Mech. 2 (15 points)

(a)

i. 3 points

For a statement that momentum is conserved or  $p_i = p_f$   
 $m v_0 = (3m) v_f$

1 point

For the correct final speed

1 point

$$v_f = \frac{v_0}{3}$$

For correct substitutions and answer

1 point

$$K_{after} = \frac{1}{2} M v^2 = \frac{1}{2} (3m) \left( \frac{v_0}{3} \right)^2 = \frac{m v_0^2}{6}$$

(1 point awarded for  $K_{after} = \frac{1}{2} (3m) v_f^2$  if student found wrong  $v_f$  or could not find  $v_f$ .)

ii. 2 points

$$\Delta K = K_{after} - K_{before} = \frac{m v_0^2}{6} - \frac{m v_0^2}{2}$$

For correct sign of answer

1 point

For correct magnitude of answer

1 point

$$\Delta K = -\frac{m v_0^2}{3}$$

(2 points awarded for any wrong answer from (a)i. minus  $\frac{1}{2} m v_0^2$ .)

(1 point awarded for  $\frac{1}{2} m v_0^2$  minus any wrong answer from (a)i.)

(b)

i. 2 points

For correct substitutions into the center of mass equation

1 point

$$r_{cm} = \frac{\sum m_i r_i}{\sum m_i} = \frac{m(0) + 2m(\ell)}{m + 2m}$$


For correct answer

1 point

$$r_{cm} = \frac{2}{3} \ell$$

Mech. 2 (continued)

ii. 1 point



For vertical arrow anywhere on diagram or in answer space

1 point

iii. 1 point

Linear momentum is conserved.

$$\mathbf{p}_i = \mathbf{p}_f$$

$$m v_0 + 3m(0) = (3m)v_f$$

For correct answer

$$v_f = \frac{v_0}{3}$$

1 point

iv. 3 points

Angular momentum is conserved.

$$L_{\text{before}} = L_{\text{after}}$$

For determining the angular momenta about the center of mass

1 point

$$L_{\text{before}} = m v_0 R \sin \theta = m v_0 \left( \frac{1}{3} \ell \right)$$

$$L_{\text{after}} = \omega I$$

For determining the moment of inertia

1 point

$$I = \sum m r^2 = m \left( \frac{2}{3} \ell \right)^2 + 2m \left( \frac{1}{3} \ell \right)^2 = \frac{2}{3} m \ell^2$$

Substituting into  $L_{\text{before}} = L_{\text{after}}$ ,

$$m v_0 \left( \frac{1}{3} \ell \right) = \frac{2}{3} m \ell^2 \omega$$

For correct answer

1 point

$$\omega = \frac{v_0}{2\ell}$$

1998 Physics C Solutions

Distribution  
of points

Mech. 2 (continued)

v. 3 points

$$K_i = \frac{1}{2} m v_0^2$$

For recognizing that final kinetic energy is translational plus rotational

1 point

$$K_f = \frac{1}{2} m v_f^2 + \frac{1}{2} I \omega^2$$

For correct substitutions and final kinetic energy

1 point

$$K_f = \frac{1}{2} (3m) \left( \frac{v_0}{3} \right)^2 + \frac{1}{2} \left( \frac{2}{3} m \ell \right) \left( \frac{v_0}{2\ell} \right)^2 = \frac{1}{4} m v_0^2$$

$$\Delta K = K_f - K_i = \frac{1}{4} m v_0^2 - \frac{1}{2} m v_0^2$$

For correct answer

1 point

$$\Delta K = -\frac{1}{4} m v_0^2$$

(Correct answer point awarded for either positive or negative sign.)

1998 Physics C Solutions

Distribution  
of points

Mech. 3 (15 points)

(a) 5 points

For each correct vector -- ½ point

For each correct magnitude -- ½ point

If the score for part (a) contained an odd number of half-points, the total score was truncated by dropping one half-point.

i.



$$N_1 = m_1 g$$

1 point


ii.



$$f_1 = 0$$

1 point


iii.



$$T = Mg$$

1 point


iv.



$$N_2 = (m_1 + m_2)g$$

1 point

v.



$$f_2 = Mg$$

1 point

(b) 3 points

For expression for the maximum frictional force

1 point

$$f_{2(\max)} = \mu_{s2} N_2 = \mu_{s2} (m_1 + m_2) g$$

For equating this force to the tension  $T = Mg$

1 point

$$Mg = \mu_{s2} (m_1 + m_2) g$$

For the correct answer


1 point

$$M = \mu_{s2} (m_1 + m_2)$$



Mech. 3 (continued)

(c) 3 points



For correctly applying Newton's second law to the hanging block

1 point

$$\Sigma F = ma$$

$$Mg - T = Ma \quad (\text{equation 1})$$

For correctly applying Newton's second law to the system of the two blocks on the plane

1 point

$$\Sigma F = (m_1 + m_2)a$$

$$T - f_2 = (m_1 + m_2)a \quad (\text{equation 2})$$

For combining equations 1 and 2 to eliminate  $T$ , substituting for  $f_2$  and solving for  $a$ 

1 point

For example, solve each equation for  $T$  and set them equal.


$$f_2 + (m_1 + m_2)a = Mg - Ma$$

$$\mu_{k2}(m_1 + m_2)g + (m_1 + m_2)a = Mg - Ma$$

$$a = \left[ \frac{M - \mu_{k2}(m_1 + m_2)}{M + m_1 + m_2} \right] g$$

(Alternate solution)

(Alternate points)



Apply Newton's second law to the three-block system, realizing that the pulley acts only to change the direction of the force produced by the tension in the string.

$$\Sigma F = m_s a$$

For correct substitutions in left side of equation above

1 point

For correct substitutions in right side of equation above

1 point

$$Mg - \mu_{k2}(m_1 + m_2)g = (M + m_1 + m_2)a$$

For correct solution for  $a$ 

1 point

$$a = \left[ \frac{M - \mu_{k2}(m_1 + m_2)}{M + m_1 + m_2} \right] g$$

Mech. 3 (continued)

(d)

i. 2 points

$$a_1 = \frac{f_1}{m_1}$$

For correct value of  $f_1$ 

$$f_1 = \mu_{k1} m_1 g$$



1 point

For correct answer

$$a_1 = \frac{\mu_{k1} m_1 g}{m_1} = \mu_{k1} g$$

1 point

ii. 2 points



Apply Newton's second law to the hanging block.

$$\Sigma F = ma$$

$$Mg - T = Ma_2 \text{ (equation 1)}$$

For correctly applying Newton's second law to block 2

1 point

$$\Sigma F = ma$$

$$T - f_1 - f_2 = m_2 a_2 \text{ (equation 2)}$$

For combining equations 1 and 2 to eliminate  $T$ , substituting for the frictional forces and solving for  $a_2$ .

1 point

For example, solve each equation for  $T$  and set them equal.

$$f_1 + f_2 + m_2 a_2 = Mg - Ma_2$$

$$Ma_2 + m_2 a_2 = Mg - \mu_{k1} m_1 g - \mu_{k2} (m_1 + m_2) g$$


$$a_2 = \left[ \frac{M - \mu_{k1} m_1 - \mu_{k2} (m_1 + m_2)}{M + m_2} \right] g$$

## 1998 Physics C Solutions

Distribution  
of points

E &amp; M 1 (15 points)

(a) 4 points

For magnitudes of electric and gravitational forces,  $F_g$  and  $F_e$ 

1 point

For equating  $F_g$  and  $F_e$  to components of  $T$ 

1 point

$$T \cos \theta = F_g = mg$$

$$T \sin \theta = F_e = \frac{kQ_A Q_B}{r^2}$$

Eliminate  $T$  between these two equations and solve for  $Q_B$ .For example from 1<sup>st</sup> equation,

$$T = \frac{(0.025 \text{ kg})(9.8 \text{ m/s}^2)}{\cos 20^\circ} = 0.26 \text{ N}$$

Substituting into the 2<sup>nd</sup> equation,

$$(0.26 \text{ N}) \sin 20^\circ = \frac{(9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2)(120 \times 10^{-6} \text{ C})Q_B}{(1.5 \text{ m})^2}$$

For proper substitutions

1 point

For the correct answer

1 point

$$Q = 1.86 \times 10^{-7} \text{ C (or } 1.9 \times 10^{-7} \text{ C)}$$

(b) 2 points

For correct answer

1 point

The new equilibrium angle will be less than  $20^\circ$  (or it decreases).For reasonable justification that indicates charges moved on conductor so positive charge is farther apart or that indicates that  $F_e$  is smaller.

1 point

Example:

The conductor allows charges to move. Positive charge will be on the far side of  $B$  from  $A$  and with a greater distance the electric force will be smaller.

## 1998 Physics C Solutions

Distribution  
of points

E &amp; M 1 (continued)

(c) 3 points

Using Gauss's law

$$\oint \mathbf{E} \cdot d\mathbf{A} = \frac{q}{\epsilon_0}$$

For evaluation of integral in equation above

1 point

For correct expression for charge enclosed

1 point

$$E(2\pi r\ell) = \frac{\lambda\ell}{\epsilon_0}$$

$$E = \frac{\lambda}{2\pi\epsilon_0} = \frac{0.10 \times 10^{-6} \text{ C/m}}{2\pi(8.85 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2)}$$

For either of the two expressions above for  $E$ 

1 point

$$E = \frac{1.8 \times 10^3}{r} \text{ N/C}$$

(d) 2 points

$$F = qE$$

For proper substitutions in equation above

1 point

$$F = (120 \times 10^{-6} \text{ C}) \left( \frac{1.8 \times 10^3}{1.5} \right) \text{ N/C}$$

For correct answer

1 point

$$F = 0.14 \text{ N}$$

(e) 4 points

For expression for work, such as  $W = q\Delta V$  or  $\int qE dr$ 

1 point

For integral with proper limits

1 point

For the proper evaluation of integral and substitution

1 point

For example,

$$W = \int_a^b \mathbf{F} \cdot d\mathbf{s} = - \int_a^b qE dr = -q \int_{1.5}^3 \frac{1.8 \times 10^3}{r} dr$$

$$W = -(120 \times 10^{-6})(1.8 \times 10^3) \int_{1.5}^{0.3} \frac{1}{r} dr = -216 [\ln r]_{1.5}^{0.3} = -216(\ln 0.3 - \ln 1.5)$$

For the correct answer

1 point

$$W = 0.35 \text{ J}$$


Alternately, one could find  $\Delta V$  using a similar integral and then use  $W = q\Delta V$ , with assignment of points similar to the above to maximum of 4 points.

1998 Physics C Solutions

Distribution  
of points

E & M 2 (15 points)

(a) 2 points



For correct placement of voltmeter

2 points

(b) 3 points

For correct application of Ohm's law

1 point

$$I = \frac{\mathcal{E}}{R_1 + R_2} = \frac{20 \text{ V}}{30 \Omega}$$

For correct value of current

1 point

$$I = \frac{2}{3} \text{ A}$$

For correct value of voltage across  $R_1$

1 point

$$V_1 = IR = \frac{2}{3} \text{ A} \times 10 \Omega = 6.67 \text{ V}$$

(Alternate solution using voltage divider)

(Alternate points)

For voltage divider equation

1 point

$$V = \frac{R_1}{R_1 + R_2} \mathcal{E}$$

For correct substitution

1 point

$$V = \frac{10 \Omega}{10 \Omega + 20 \Omega} \times 20 \text{ V}$$

For correct answer

1 point

$$V = 6.67 \text{ V}$$

(c)

i. 2 points

For correct answer

2 points

$$V = 0$$

(1 point awarded for stating that no current flows)

ii. 1 point

$$Q = CV$$

For correct substitutions and answer

1 point

$$Q = 15 \mu\text{F} \times 20 \text{ V} = 300 \mu\text{C}$$

## E &amp; M 2 (continued)

(d) 2 points

For correct answer

$$V = 0$$

2 points

(1 point awarded for no current or realization of new initial conditions)

(e)

i. 2 points

For correct application of Ohm's law and answer

2 points

$$I = \frac{\mathcal{E}}{R_1 + R_2} = \frac{20 \text{ V}}{30 \Omega} = \frac{2}{3} \text{ A}$$

(1 point awarded for  $\frac{20 \text{ V}}{10 \Omega}$ )

ii. 1 point

For correct substitution in energy equation and correct answer

1 point

$$U_L = \frac{1}{2} LI^2 = \frac{1}{2} (2 \text{ H}) \left( \frac{2}{3} \text{ A} \right)^2 = 0.444 \text{ J}$$

(1 point also awarded if incorrect current in (e)i. (except zero) was substituted and answer was consistent with this current.)

(f) 2 points

For correct equation

2 points

$$\mathcal{E} - I(R_1 + R_2) - L \frac{dI}{dt} = 0$$

(1 point only awarded if one sign incorrect)

One point was subtracted from the final score for two or more wrong or absent units for parts where the answer was given, except where the answer was zero, in which case units were not counted.

## 1998 Physics C Solutions

Distribution  
of points

E &amp; M 3 (15 points)

(a) 4 points

Using Newton's 2<sup>nd</sup> law along the ramp,

$$F_g \sin \theta - F_m = ma, \text{ where } F_g = \text{force of gravity, and } F_m = \text{magnetic force}$$

At constant speed,  $a = 0$ , so

$$F_g \sin \theta - F_m = 0$$

For the following expression for  $F_m$ 

$$F_m = F_g \sin \theta$$

1 point

For substituting  $F_g = mg$ 

$$F_m = mg \sin \theta$$

1 point

For substituting  $F_m = I\ell B$ 

$$I\ell B = mg \sin \theta$$

1 point

For correct answer

$$I = \frac{mg \sin \theta}{\ell B}$$

1 point

(b) 4 points

$$|\mathcal{E}| = \left| -\frac{d\phi_M}{dt} \right|$$

For correct expression for  $\phi_M$ 

$$\phi_M = B\ell x$$

1 point

For the time derivative of  $\phi_M$ 

$$\frac{d\phi_M}{dt} = B\ell v$$

1 point

For correct use of Ohm's law to find current

$$\mathcal{E} = IR$$

1 point

$$I = \frac{\mathcal{E}}{R} = \frac{B\ell v}{R}$$

For equating this expression to the expression for  $I$  from part (a) and solving for  $v$ 

1 point

$$\frac{B\ell v}{R} = \frac{mg \sin \theta}{B\ell}$$

$$v = \frac{mgR \sin \theta}{B^2 \ell^2}$$

## 1998 Physics C Solutions

Distribution  
of points

E &amp; M 3 (continued)

(c) 2 points

For correct expression for power

$$P = I^2 R$$

1 point

For correction substitution of the expression for  $I$  from part (a)

$$P = \frac{(m^2 g^2 \sin^2 \theta) R}{B^2 \ell^2}$$

1 point

(d) 3 points

For either of the following two expressions

$$m \frac{dv}{dt} = mg \sin \theta - I \ell B$$

$$\frac{dv}{dt} = g \sin \theta - \frac{B^2 \ell^2 v}{mR}$$

1 point

For rearranging terms in either expression to set up integral

For example,

$$\frac{dv}{g \sin \theta - \frac{B^2 \ell^2 v}{mR}} = dt$$

1 point

For integrating this expression, and applying  $v(0) = 0$  to get answer

$$v(t) = \frac{mRg \sin \theta}{B^2 \ell^2} \left[ 1 - \exp\left(-\frac{B^2 \ell^2 t}{mR}\right) \right]$$

1 point

(e) 2 points

For correct answer

Yes, the final speed of the bar decreases

1 point

For any reasonable justification

For example;

Since the two resistors are in parallel across the emf in the bar, the new effective

resistance is  $\frac{R}{2}$ . And since  $v = \frac{mgR \sin \theta}{B^2 \ell^2}$  from part (b), then if  $R$  decreases,

the speed decreases.

1 point